

Analysis of observed and perceived climate change and variability in Arsi Negele District, Ethiopia

Zenebe Mekonnen¹ · Habtemariam Kassa² · Teshale Woldeamanuel¹ · Zebene Asfaw¹

Received: 8 July 2016 / Accepted: 16 March 2017
© Springer Science+Business Media Dordrecht 2017

Abstract Climate change and variability has been detected in Ethiopia. Smallholder and subsistence farmers, pastoralists and forest-dependent households are the most hit by climate-related hazards. They have to have perception of climate change in order to respond it through making coping and/or adaptation strategies. Local perceptions and coping strategies provide a crucial foundation for community-based climate change adaptation measures. This study was specifically designed to (1) assess households' perception and knowledge in climate change and/or variability, and (2) establish the observed changes in climate parameters with community perceptions and climate anomalies. Purposive stratified random sampling method has been used to gather information from 355 sample households for individual interviews supplemented by group discussion and key informant interviews. The analysis of observed and satellite climate data for the study district showed that mean maximum and minimum temperature for the period 1983–2014 has increased by 0.047 and 0.028 °C/year, respectively. However, the total rainfall has declined by 10.16 mm per annum. Seasonally, the rainfall has declined by 2.198, 4.541, 1.814 and 1.608 mm per annum for Ethiopian summer, spring, autumn and winter seasons,

Electronic supplementary material The online version of this article (doi:[10.1007/s10668-017-9934-8](https://doi.org/10.1007/s10668-017-9934-8)) contains supplementary material, which is available to authorized users.

✉ Zenebe Mekonnen
zenebemg2014@gmail.com

Habtemariam Kassa
h.kassa@cgiar.org

Teshale Woldeamanuel
twoldeamanuel@yahoo.com

Zebene Asfaw
zebeneasfaw@gmail.com

¹ Wondo Genet College of Forestry and Natural Resources, Hawassa University, P.O. Box 128, Shashemene, Ethiopia

² Forests and Livelihoods Research, Ethiopia Office, Centre for International Forest Research (CIFOR), P.O. Box 5689, Addis Ababa, Ethiopia

respectively. Similarly, the mean maximum temperature of the study area had showed an increment of 0.035, 0.049, 0.044 and 0.065 °C per year for spring, winter, autumn and summer seasons, respectively. The observed climate variation has been confirmed by people's perception. Considering what had been the existed situations before 30 years ago as normal, an increase in temperature, an increase in drought frequency, a decrease in total rainfall, erratic nature of its distribution and the tardiness of its onset had been perceived by 88, 70, 97, 80 and 94% of the respondents, respectively, at current time—2015. Deforestation as a casual factor of climate change and variability had been perceived by 99.7% of the respondents. This had been also confirmed by scientific studies as it emits carbon dioxide and is the main driver of climate change and variability. Indigenous knowledge, including climate predictions, has been used by people to implement their day-to-day agricultural activities. Therefore, science should be integrated with the perception and indigenous knowledge of people to come up with concrete solution for climate change and variability impacts on human livelihoods.

Keywords Climate change · Climate variation · Ethiopia · Indigenous knowledge · Perception

1 Introduction

Climate change and variability is a global concern (IPCC 2014), and it has been detected in Ethiopia (NMA 2006; McSweeney et al. 2008). Climate variability refers to variations in the mean state and other statistics of the climate on both temporal and spatial scales beyond the individual weather events (IPCC 2014). While climate change is a change in climate over comparable time periods (a decade or more) which is attributed directly or indirectly to human activity that alters the composition of the global atmosphere (IPCC 2014). Between 1951 and 2006, the annual minimum temperature in Ethiopia was increased by about 0.37 °C per decade (NMA 2006; Adem and Bewket 2011). The IPCC's mid-range emission scenario (IPCC 2007) shows that the mean annual temperature in Ethiopia was increased by 1.3 °C between 1960 and 2006, which is an average rate of 0.28 °C per decade (McSweeney et al. 2008). According to this emission scenario, the mean annual temperature also expected to increase in the range of 0.9–1.1 °C by the year 2030 and 1.7–2.1 °C by the year 2050 as compared to the 1961–1990. With respect to annual precipitation, for the same scenario and base period, the expected variation is in the range of 0.6–4.9% by 2030 and 1.1–18.2% by 2050 (Mekonnen and Hailemariam 2000; NMA 2006; Regassa et al. 2010). The frequency of drought in Ethiopia particularly in the recent decades is an indication of the prevalence of the variation in climate. There were 19 drought events which occurred in Ethiopia between the period 1900 and 2002, which is almost once in 6 years in 1900–1987(14 drought events) and roughly in 3 years in 1988–2002 (five drought events) (NMSA 1987; World Bank 2005). In the period between 1999 and 2008, drought and flooding affected nearly 21 million people (19.7 million by drought and the remaining by flooding) in Ethiopia (OFDA/CRED 2009).

The livelihoods of smallholder farmers are mainly dependent on climate change and variability sensitive sectors such as agriculture and natural resources. These communities live with a range of livelihood risks (Gandure et al. 2013). Climate change impact is more intense when it is combined with other environmental changes (Mertz et al. 2009; IPCC 2014). Households that perceived the variability and change in climate can cope and adapt

to the change, hence less vulnerable to the impacts than those who did not perceive it (UNFCCC 2007). These households respond to the change and variability via practicing different indigenous coping and adaptation techniques as the perception enables their decision-making (Gyampoh et al. 2009; NAC 2010; Osbahr et al. 2011; Kumar 2014; Tilahun and Bedemo 2014; Huda et al. 2016; IFAD 2016).

Hence, perception of the long-term changes in climate by them plays an important role in shaping their behaviour (Abid et al. 2014). Perceptions of risk and cognitive process of primary decision makers are important for the adoption of different adaptation decisions (Grothmann and Patt 2005; Nyanga et al. 2011; Simelton et al. 2011; Amdu et al. 2013; Burnham 2014; Berman *et.al.* 2014). In this regard, several studies (e.g. Egeru 2012; Amdu et al. 2013; Varadan and Kumar 2014) indicated smallholder farmers' experience of climate change and variability regarding decreasing and unpredictable rainfall, increasing temperature and delayed onset of rainfall, which were also validated with actual climate trend analysis. As smallholder farmers are engaged actively with their natural environment in their day-to-day lives, they have accumulated sizable and sophisticated bodies of knowledge and practices about their environment, its variability and transformation which is a basis for indigenous adaptation and provide important insights into processes of modern adaptation to climate variability and change (Nakashima et al. 2012). In our context, indigenous, local or community and traditional knowledge are used synonymously. Community-based and local knowledge may offer valuable insights into environmental change instigated by climate change and complement broader-scale scientific research with local precision and nuance. Ignorance of such knowledge, however, reduces individual perception of the risks posed by climate change as well as affects the costs and benefits of different adaptation options: adjustment in natural or human systems in response to actual or expected climatic stimuli or their effects, which moderates harm or exploits beneficial opportunities (IPCC 2014). Local perceptions and knowledge provide a crucial foundation for community-based climate change adaptation measures (McLean et al. 2011; Nakashima et al. 2012; Ramos and McLean 2012) and are acknowledged by IPCC as invaluable for developing adaptation and natural resource management strategies in response to environmental and other forms of change (IPCC 2007; Gyampoh et al. 2009; The Anchorage Declaration 2009; IPCC 2010a; IPCC 2013; UNFCCC 2010; Magni 2016; Ford et al. 2016). As the local perceptions and the resulting indigenous knowledge are location and site specific, there investigation in different socioeconomic setting is worthwhile. Nevertheless, limited attention has been given to the knowledge and practice emanated from the perception of smallholder farmers and the accumulated knowledge and long-time experience about their environment. The knowledge and practice in this regard are very important as the combination of the knowledge obtained from the community's perceptions and observed data can be used for better understanding about climate change and variability in general and what response strategies can be taken. This study was specifically designed to (1) assess households' perception and knowledge of climate change and/or variability and (2) establish the observed changes in climate parameters with community perceptions and climate anomalies.

2 Materials and methods

2.1 Description of the study area

The study was conducted in Arsi Negele District, which is located in West Arsi zone of the Oromia Regional State, Ethiopia. The district is located between 7.15 and 7.75°N latitudes

and 38.35 and 38.95°E longitudes (Fig. 1). The altitude of the district ranges between 1500 and 3000 m above sea level. The topography encompassed the central rift valley floor, including lakes Langano, Shalla and Abijata, and extended to the eastern escarpment of the central rift valley of Ethiopia. The district's average annual temperature varies between 10 and 25 °C while the annual rainfall varies between 500 and 1000 mm (Fig. 2).

Demographically, between the periods 1994 and 2007, the rural population was increased by 83% while the urban population increased by 119%. The rural population of the district accounts 83% of the total population in 1994 and 80% in 2007 (ENC 1994, 2007; CSA 2005).

3 Methods

In this study, both the qualitative and quantitative data collection methods were employed. The study was a cross-sectional survey of respondents from six kebeles, the lowest administrative division in Ethiopia's geopolitical administration, and three agro-ecologies—two kebeles from each agro-ecology. This was supplemented with 18 focus group discussions, three from each kebele (2 kebeles/ACZ × 3 focus groups/kebele × 3 ACZs = 18 focus groups), and 30 key informant interviews, five from each kebele. It also complemented by secondary sources from measured and satellite climatic data.

3.1 Sampling design

This study employed multistage sampling techniques. In the first stage, Arsi Negele District was selected purposively as it represents three distinct agro-ecologies such as lowland, midland and highland, which can be used to show the spatial dimensions of climate change and variability. In the second stage, six kebeles, two each from each agro-ecology, were




Fig. 1 Map of the study area




Fig. 2 Diagram showing variation in climate variables over Arsi Negele District (1983–2014)

selected based on the representativeness of an agro-ecology. The third stage was selecting the sample households from 4257 households in the area. A total of 355 households (163 poor, 124 medium and 68 better off) were selected through stratified random sampling. Then proportional numbers of households at each kebele were randomly selected based on wealth categories and gender for household interview. The sample representing the population was selected on the basis of normal distribution with confidence level of 95% and margin of error 5% followed by finite population correction (FPC) to get the true sample (i.e. $n = (s \times N)/(s + N - 1)$, where n is true sample and s is sample of 50% distribution ~ 385 , $N = 4257$) (Israel 1992; Bartlett et al. 2001).

Hence, to collect data at household and community level, three major primary data collection methods were used. These include key informant interviews, focus group discussion and household survey.

As key informants interview, those people who have relatively knowledgeable about their community situation, local natural resources, climate conditions, the culture of the community, overall development and the respective changes in these are used (Smith and Sharp 2012). Hence, they shared their built-up knowledge and experience to the interviewer. To select these key informants, a snowball method at which one key informant was contacted based on kebele officials information and document has been used. Then, he/she would inform us the second, the second would tell the third and so forth until a saturation number was reached (Bernard 2006).

The focus group discussions were made representative of the community including youth, women and elders. These focus group discussions were made with three independent groups per kebele at which each group consisted of 8–10 peoples. The focus group discussions were on different issues including farming calendar, establishing local criteria for wealth categorization, perception of climate change and variability (temperature and rainfall) as well as environmental degradation and causes of local climate variation. The members of a group were selected randomly including women, elders and youth to obtain varied knowledge and views. At least one key informant was included in each group to triangulate information. The focus group discussion had made use of participants' feelings, perceptions and opinion about local climate variations and change. To understand climate change and variability in local context, we, the facilitator and FGD participants, first have defined what was considered normal in local context before 30 years ago.

The household survey was conducted with sample households. To select the sample households, list of all households has been taken from kebele administration office which were already categorized into three groups, namely poor, medium and better off—based on the kebele's wealth categorization criteria. After this, we had discussed on the list with key

informants in the community with those categories and had allowed them to categorize the households with their own criterion such as asset possession, better house, income and fertile farmland. Those that have intersected with the kebele and key informants' categorizations were considered in the sampling randomization. Then, proportional number from each category has been taken randomly—163 poor, 124 medium and 68 better off—making a total of 355 sample households. Sample households were taken from each wealth category proportionally to improve the reliability of the data and to capture differential perception and knowledge about climate change and variability among social status. The interview was designed in semi-structured and structured forms and translated into local language—*Afan Oromo*.

As secondary data, satellite images and meteorological data obtained from National Meteorological Agency (NMA) were used for observed climate trend analysis. Using the satellite *data*, Arsi Negele District has been divided into 49 grids cells of size 10×10 km spatial and daily/monthly temporal resolutions, and climatic data have been reconstructed by NMA personnel in 2015 for the geographic location of the district between 7.15 and 7.75°N and 38.35 and 39.95°E . Then, latitude lines 7.25 , 7.35 and 7.55°N were chosen representing and dissecting the lowland, midland and highland agro-ecologies of the district, respectively.

4 Data analysis

Data were analysed using descriptive statistics such as frequencies, means, percentages and graphs as well as time series analysis using ordinary least square for temperature and rainfall. In addition, correlational statistics are used to describe the relationship and prediction of climate variables: observed vs. perceived. Statistical software such as SPSS v. 20 and Minitab v. 17 was used to analyse the quantitative data obtained from survey and secondary information (meteorological data). The qualitative data from focus group discussions and key informants interviews were analysed and interpreted in the form of narratives, trend analysis and descriptions. The reconstructed grid cells climate data were compared with observed climate data from weather station. Comparison was made for weather station located in the district, and an F-test was made for those years with full temperature and rainfall data to see whether there is statistical significant difference between the mean annual values of the observed and gridded data types. The F-test analysis showed that there is no statistical significant difference between the two data sets. As a result, satellite data were used for temporal and spatial climate variability analysis which avoids the missed data gaps in the observed temperature and rainfall data (Dinku et al. 2013, 2014).

5 Results

5.1 Observed climate variability

The analysis of observed climate data shows that there was variability in climate in Arsi Negele District. This is supported by the analysis of both temperature and rainfall data. The result shows that the variation has both temporal and spatial dimension. The analysis of temperature data from 1983 to 2014 showed that the mean maximum temperature varies

from 24.52 °C in 1993 to 27.76 °C in 2009. The ranges of variation for the mean annual maximum, minimum and average temperatures were $[-2.28, 2.42]$, $[-1.72, 3.82]$ and $[-1.48, 3.08]$ standard deviations, respectively (Fig. 3b). As indicated in Fig. 3a from Arsi Negele Station (7.35°N, 38.66°E, 1913 m a.s.l.), the mean annual maximum temperature, the mean annual minimum temperature and mean annual temperature showed an increasing trend of 0.047, 0.028 and 0.038 °C per annum, respectively (Fig. 3a). For the mean maximum temperature in the study area, the maximum positive deviations were observed in the years 1988, 1999 and 2012 and the maximum negative deviation in 2010. Similarly, the maximum positive deviation for the minimum temperature was observed in 1988 while the negative highest deviation was in 2014. For the mean annual temperature, the highest positive and negative deviations were in 2012 and 2000, respectively. The rainfall data analysis for the period 1983–2014 has showed that there was variation in the total annual rainfall (TARF) with the peak in 2004 with 1486.5 mm and sharp drop in 2013 with 206 mm. This variation ranges from -2.44 to 2.42 standard deviations from the mean for Arsi Negele Station (Fig. 3b). The total annual rainfall has declined by 10.16 mm per annum with the highest positive and negative deviations in 2003 and 2013, respectively.

5.2 Spatial and temporal variations

The analysis of satellite data for the period 1982–2011 has showed that there was variation in temperature and rainfall both spatially and temporally. For the grid cells along 7.15°N latitude, the temperature and rainfall showed decreasing trend as one goes from west to east except that the minimum temperature was at peak for the grid cell 38.65°E in both periods (Fig. 4i). The minimum temperature along the 7.35°N grid cells showed somewhat declining trend at first, then increasing from 38.45 to 38.75°E and then somewhat decline when running from west to east for both the three decades (Fig. 4d). While the maximum temperature has showed a general declining trend for both periods (Fig. 4e), the rainfall has showed a declining trend for the grid cells between 38.35 and 38.55°E. It continued with similar trend for the decades 1992–2001 and 2002–2011 and increasing trend for the decade 1982–1991 (Fig. 4f) as one goes from east to west. For the grid cells along 7.55°N which represents the lowland climate, the temperature has showed an increasing trend and then somewhat declined.




Fig. 3 Temperature and rainfall trends (a) and their respective standardized anomalies (b) from Arsi Negele Station




Fig. 4 Spatial and temporal climate variability in Arsi Negele District

The rainfall has generally showed a declining trend for both decades as we go from east to west (Fig. 4g–i). The analysis on a same grid cell showed that the temperature was more or less higher for the decade 1982–1991 than for the other two decades, but the average temperature for 2002–2011 is higher than 1992–2001 for all grid cells along 7.25, 7.35 and 7.55°N. Nevertheless, the total annual rainfall was generally lower in 2002–2011 than in 1982–1991 and 1992–2001 at almost all of the grid cells along latitude 7.25 and 7.35°N. However, the total rainfall was higher for the decade 2002–2011 for the grid cells along 7.55°N latitude (Fig. 4a–i).

5.3 Seasonal climatic variations

5.3.1 Rainfall

The analysis of climatic variables in the study area showed a high variability of rainfall with r^2 ranging from 0.023 for June–July–August (JJA)—Ethiopian summer season—to 0.090 for March–April–May (MAM)—Ethiopian spring season—during the period 1983/1984–2014/2015 (Fig. 5). During this same period, the rainfall has declined by 2.198,

4.541, 1.814 and 1.608 mm per annum for JJA, MAM, September–October–November (SON)—Ethiopian autumn season—and December–January–February (DJF)—Ethiopian winter seasons—respectively.

5.3.2 Minimum temperature

The variation in minimum temperature in the study area is high. The r^2 values for all four seasons are very small, indicating that there is high variation. In the study area, the mean minimum temperature has showed a decrease of 0.014 and 0.016 °C per year for summer and spring seasons, respectively. On the other hand, the minimum temperature has showed an increase of 0.07 and 0.069 °C per annum for autumn and winter seasons, respectively (Fig. 6).

5.3.3 Maximum temperature

The analysis for the maximum temperature in the study area showed the highest variability for spring season with $r^2 = 0.096$ followed by autumn ($r^2 = 0.187$), winter ($r^2 = 0.217$) and summer ($r^2 = 0.317$), respectively. The mean maximum temperature of the study area had showed an increment of 0.035, 0.049, 0.044 and 0.065 °C per year for spring, winter, autumn and summer seasons, respectively (Fig. 7).

5.4 Perceptions of climate variability

Households and communities perception to climate change and variability in the study area was also assessed using focus group discussions and key informants interviews. In this regard, both sources underlined that there was clearly climate change and variability in their village as compared to 30 years ago. They perceived this change in terms of variation in temperature and rainfall. They characterized the variation in rainfall as declined in amount, discontinuous in distribution and erratic in its onset and ending. Regarding the variation in temperature, it was indicated as it has been increased to the extent they felt it as they could not wear clothes which they used to wear before. The respondents explained the




Fig. 5 Seasonal variation in rainfall in Arsi Negele District (where x is year 1, 2, 3... 32 at which 1 is year 1983 and 32 is year 2014)




Fig. 6 Seasonal variation in minimum temperature from Arsi Negele District (where x is year 1, 2, 3... 32 at which 1 is year 1983 and 32 is year 2014)




Fig. 7 Seasonal variation in maximum temperature from Arsi Negele District (where x is year 1, 2, 3... 32 at which 1 is year 1983 and 32 is year 2014)

variation in climate change with respect to three different periods. These periods were the periods in which three different regimes in Ethiopia existed. The first period was the period called Imperial (pre-1974), the second period was when the Dergue (Military committee) government took power (1974–1991), and the third period is the period the incumbent regime/EPRDF (since 1991). In this regard, they indicated that the climate was better during the Imperial than the Dergue and was better during the Dergue than the present, they added. Even in the year 1984, which was called the drought year in Ethiopian drought

history, there was slight rain in summer, but nowadays the drought and intensity of heat have become more severe than that, the witnesses supplemented. The key informants and focus group members asserted that the rain season was longer and its distribution was uniform in the past than nowadays.

The responses of group discussion and key informants were similarly reflected in individual household interview (Table 1). These farmers' perception of increasing temperature and declining rainfall is similar to the observed trends discussed in the previous sections except that farmers give more focus on the distribution of rainfall rather than the total amount that can be downpour at once. Besides, people in group discussions have described that a small rain for continuous 10 days is better for crop growth than a heavy rain for only 1 or 2 days.

Farmers have recognized the variations and changes in climate by cognitive learning, awareness given by different communication tools as well as from indigenous knowledge. For example, about 98.4% of respondents perceived the existence of climate change and variability by feeling warm due to increased temperature than usual (Table 2).

Table 1 Trends of climate and climate-related variables as perceived by farmers

Climate/related variables	Scales compared to 30 years ago	Percent of respondents who realized the variations			
		Lowland (<i>n</i> = 104)	Midland (<i>n</i> = 103)	Highland (<i>n</i> = 148)	Total (<i>N</i> = 355)
Total rainfall	No change	0.00	0.00	1.67	0.63
	Increased	2.00	0.00	4.17	2.19
	Decreased	98.00	100.00	94.17	97.19
Rainfall distribution	No change	0.00	0.00	7.50	2.81
	Uniform	7.00	20.00	24.17	17.50
	Erratic	93.00	80.00	68.33	79.69
Rainfall onset	No change	0.00	0.00	0.83	0.31
	Delayed	97.00	100.00	87.50	94.38
	Early	3.00	0.00	11.67	5.31
Average temperature	No change	0.00	0.00	1.67	0.63
	Increased	95.00	95.00	77.50	88.44
	Decreased	5.00	5.00	20.83	10.94
Frost occurrence	No change	1.00	5.00	12.50	6.56
	Increased	49.00	37.00	31.67	38.75
	Decreased	50.00	58.00	55.83	54.69
Hailstorm frequency	No change	9.00	7.00	25.00	14.38
	Increased	38.00	35.00	18.33	29.69
	Decreased	53.00	58.00	56.67	55.94
Drought frequency	No change	0.00	1.00	11.67	4.69
	Increased	79.00	70.00	61.67	69.69
	Decreased	21.00	29.00	26.67	25.63
Flood frequency	No change	3.00	0.00	8.33	4.06
	Increased	34.00	47.00	61.67	48.44
	Decreased	63.00	53.00	30.00	47.50

Table 2 Different means of knowing the existence of climate change and variability by respondents

Means of knowing	Percent of respondents answering yes				Chi-square (χ^2)
	Lowland ($n = 104$)	Midland ($n = 103$)	Highland ($n = 148$)	Total ($N = 355$)	
Learned from cognition					
Observing changes in tree flowering	87.00	92.00	86.64	88.40	189.113a
Observing changes in wind speed and frequency	89.00	93.00	87.49	89.70	201.613a
Observing fluctuation in total rainfall	93.00	94.00	96.66	94.70	255.613a
Observing erratic and unpredictable rainfall	98.00	99.00	93.33	96.60	277.513a
Observing changes in animal behaviour	96.00	87.00	91.68	91.60	221.113a
Feeling warm due to increased temperature	100.00	100.00	95.84	98.40	300.313a
Observing changes in the timing of rainfall onset	72.00	77.00	70.82	73.10	68.450a
Learned from different communications tools					
Weather forecast and report form TV news	29.00	39.00	61.64	44.40	4.050a
Weather forecast and report form radio news	78.00	75.00	87.52	80.60	120.050a
Extension service provided by public organizations	84.00	97.00	88.33	89.70	201.613a
Extension services by civic organizations	75.00	98.00	81.64	84.70	154.013a
Public meetings	98.00	98.00	95.83	97.20	285.013a
Development committee and/or 1:5 farmers arrangement	95.00	97.00	92.48	94.70	255.613a
Field demonstration on climate impacts	86.00	92.00	86.67	88.10	186.050a
Public education on climate change	78.00	90.00	80.83	82.80	137.813a

(a) No cells have expected frequencies less than 5. The minimum expected cell frequency is 178 for the total

5.4.1 Indigenous knowledge about climate

The role indigenous knowledge with respect to climate variation and change prediction is known to be very important in the area. That is, the communities in the study area, especially elders and local *Aba Geda*—traditional leader of the Oromo community—have accustomed to several indigenous knowledges to predict climatic situations (Table 3). These communities indigenous knowledges (CIK) play important role, not less than the information from National Meteorology Agency (Woldekiros Dube, 07/12/2015, Key informant, among others), in that farmers could adjust, plan and implement their farming activities accordingly. Key informants dictated that the main growing season in the study

Table 3 Community indigenous knowledge and their local narratives of predicting climate

Bases of CIK	Local narratives and interpretation
Size and position of the moon	When the size of the moon is full and tilted to the left, we believe that there will be good rainfall When its size becomes very thin (crescent) and has no tilt in its position, we believe that there will be no rainfall
Length of dry season	When the number of dry months is five or more in a year, we considered it a drought year When the number of dry months is four or less in a year, we considered it a good rainy year
Wind direction	When the wind starts to blow from highland to lowland, we believe that the rain is to come soon in the next few days When the wind starts to blow from lowland to highland, we believe that there will be no rain in the locality
Cloud speed and direction	When the cloud moves fastest from west to east in a particular day, we believe as it rains just on that day When clouds move slower from west to east in a particular day, we believe as it rains the next day When clouds change direction from normal, we believe as it results in no rain
Timing of rainbow	When rainbow is observed after rain, we believe as the rain is going to stop When the rainbow is observed at the period of no rain, we believe as the rain is going to come
Timing of rainfall	When the rain start to come during sunset, we believe as it is “good summer” with enough rainfall When the rain start to come during sunrise, we believe as it is “poor summer” with shortage of rainfall
Animal behaviour	When bulls started to caper from here to there in the field in a dry time, we believe that the rain is going to come soon Bees will migrate (“godinsa kanissa”) to the lowlands (Langano area) when the rain starts to rain in that area and come back to the highlands when it stops When red colour ants start to roam around people’s home and around fences, we consider it as indicator that rain is going to come When hyenas scream (“warabeesii yoo yusuu”) in a slow and continuous ton and roam around village and break fences to eat what it gets, we considered as the rain is going to come
Tree flowering	Most of the trees including acacias, ficus spp., <i>Croton macrostachyus</i> (Del.), <i>Celtis africana</i> (Burm.) and <i>Podocarpus falcatus</i> (Thunb.) have shifted their flowering period at least by 1 month earlier or delayed Most of the above species produce immature seeds even after flowing due to the shortage of rainfall distribution, they said

district is June to August (summer in Ethiopia). The normal sowing dates for xafii (Orm), *teff* (Amh.), *Eragrostis tef* (Zucc.) are end of July to beginning of August. In the beginning of the 2015 growing season, there were predictions by CIK and NMA in reference to the prevailing El Nino events. On the one hand, key informants claim that by observing the size and position of the moon they predicted as the rain will stop in September and gave advice to make adjustment on the sowing dates of *E.tef* to the middle of July. On the other hand, farmers heard through radio that NMA has predicted as the rain will extended up to November and advised farmers to extend the sowing dates of *E.tef* at the end of August. Finally, the rain did stop in September as predicted by CIK while most farmers had sown

E.tef as suggested by NMA and they lost production due to shortage of rain, key informants said. This shows that lack of integration of CIK and scientific knowledge leads to misconception on farmers' agricultural activities.

The change in cropping calendar is indicated as one of the evidences for climate change and variability in the study area. In the study area, the main stay of livelihood in the study area is mixed agriculture at which crop production takes the lion share. However, crop cultivation is mostly rain fed and highly impacted by rainfall distribution. In the focus group discussions, the farmers explained that the delay on rainfall onset has obliged them to shift their cropping calendar what they were accustomed before 30 years ago (1985) (Tables 4, 5). It has been understood from farmers' opinions that some crops have shifted their agronomic ranges towards the highland and lowland agro-climatic zones. Wheat is the case in point which did not grow in the lowlands and higher highlands before 30 years ago from 2015, while it does at the current times.

5.4.2 Perception on causes of climate change and variability

Regarding causes of climate variability, the respondents suggested that the rainfall decline followed the decline in deforestation/forest cover and the decline in forest cover followed the increase in population and agricultural expansion; the temperature has increased due to the shift of warmer (lowland) climate into the highlands; industrial expansion in nearby towns and other areas as we learnt from extension and radio, but do not know the scientific bases; and the other cause is, as we believe, is natural phenomena (Fig. 8).

6 Discussion

6.1 Observed climate change

Climate change is assessed by using meteorological observations and/or farmers perceptions. The analysis of climate variables from the reconstructed grid cells and actual measurements for the study district has showed that there is high variability of temperature and rainfall both spatially and temporally. This variability has been proved for other parts of Ethiopia and Africa (McSweeney et al. 2008; Adem and Bewket 2011; Okonya et al. 2013; Amekudzi et al. 2015; Asante and Amuakwa-Mensah 2015; Tierney et al. 2015; McPhaden 2015; Muluneh et al. 2016). The rainfall variability in Ethiopia is generally correlated with the movement of the Inter-Tropical Convergence Zone (ITCZ) in the long rain season and with the warming of the Indian Ocean in the short rain season (Simelton et al. 2011). For the study area in general, the temperature has showed an increasing trend while the rainfall has showed a declining trend. Observed climate parameters from stations which are in the adjacent districts (lowland) to the study district showed that the rainfall varies between $[-1.46, 2.5]$, $[-2.36, 3.0]$ and $[-1.50, 2.0]$ standard deviations from the mean for Bulbula (7.72°N, 38.65°E, 1606 m a.s.l.), Ziway (7.933°N, 38.7°E, 1640 m a.s.l.) and Adami Tulu (7.856°N, 38.703°E, 1653 m a.s.l.) stations, respectively (Fig. 9). Additional climate data from Degaga station (7.43°N, 38.84°E, 2067 m a.s.l.) which reside in the study district (highland) also showed similar trend $[-1.83, 2.15]$.

Table 4 The Cropping calendar of the major crops in lowland areas

Rainfall & Activities	Time scale	Months											
		J	F	M	A	M	J	J	A	S	O	N	D
Rainfall onset and ending	30 2015		[Dark Blue Bar]					[Dark Blue Bar]					
Sowing	Maize	30 2015			[Dark Blue]								
	Haricot bean	30 2015				[Orange]							
	Tef	30 2015					[Dark Blue]						
	Finger Millet	30 2015				[Orange]							
	wheat	30 2015											
	Barley	30 2015											
	Did not grow												
	Did not grow												
Did not grow													
Harvesting	Maize	30 2015										[Dark Blue]	
	Haricot bean	30 2015										[Dark Blue]	
	tef	30 2015										[Dark Blue]	
	Finger Millet	30 2015										[Orange]	
	wheat	30 2015											[Orange]
	Barley	30 2015											[Orange]
	Did not grow												
	Did not grow												

6.2 Perceived climate change

The analysis of the household survey, key informant interviews and focus group discussions shows the reality of climate change and variability in their locality. Considering what




Fig. 8 Causes of local climate change and variability as perceived by farmers




Fig. 9 Standardized rainfall anomalies at different stations within and nearby Arsi Negele District

respondents, respectively. This was in line with the observed climate trend. The perceived delayed onset of rainfall is supported by meteorological evidence of declining rainfall, erratic rainfall distribution by seasonal variability, rainfall termination by declining monthly rainfall and shorter rainfall duration by increase in number of dry days (Simelton et al. 2011). Deforestation as a casual factor of climate change and variability had been perceived by 99.7% of the respondents. This anthropogenic activity had been also confirmed by scientific studies as it emits CO₂ and is the main driver of climate change and variability (Houghton et al. 2012; FAO 2014; IPCC 2014; Stocker and Joos 2015). Studies

in different parts of the world underscored that indigenous knowledge, including climate predictions, has been used by people to implement their day-to-day agricultural activities (Egeru 2012; Moyo et al. 2012; Varadan and Kumar 2014; Ndambiri et al. 2014). Farmers also use an adjustment to their agricultural calendar as a response to climate variability (Yegbemey et al. 2014). Simelton et al. (2011) stated that this knowledge of farmers' perceptions of how climate is changing is crucial in anticipating the effects of climate change, as only farmers who perceive a problem will adapt to it, indicating that assessing the impact of erratic rainfall, for example, is more important for farmers' decision and outcomes than crop models with meteorological data. In this study, it was understood that indigenous knowledge and perception of climate change and variability by farmers have enabled them to devise local coping strategies that are oriented towards survival such as crop diversification and firewood sale to escape from immediate climate change impacts. These local-level coping strategies are the basis for adaptation to climate change which again enables them to devise adaptation strategies that are oriented towards longer-term livelihoods security (Dazé et al. 2009) such as water harvesting, enhancement in irrigation, mulching, changing sowing date and time of soil cultivation, soil fertility management, minimized post-harvest losses.

6.3 Strengths and limitations of observed, satellite and perceived climate change and variability

Observed climate change and variability analysis has its strength by being used as a basis for satellite climate data, more focus on meteorological drought (based on below average annual rainfall) and has got standard and is more reliable. It has limitations of incomplete data and less focus on agronomic drought (due to increase in temperature and evaporation) (Simelton et al. 2011). Climate change and variability analysis from satellite data provides information on temperature and precipitation even over remote parts where weather stations are not available. This is very useful to fill gaps in the time series at which weather stations suffered from. In this study, the correlations between satellite data and measured data were highly significant for rainfall ($p < 0.001$, $r^2 = 0.91$), minimum temperature ($p < 0.001$, $r^2 = 0.99$) and maximum temperature ($p < 0.001$, $r^2 = 0.95$). Despite its usefulness, satellite climate data have shortcomings: (1) time period of observation is very short, (2) accuracy at higher temporal and spatial resolutions might be poor, and (3) it lacks homogeneity in time series (Dinku et al. 2013). In parallel, downscaled climate data analysis uses complex mathematical rules and helps to predict future trends even if it has limited use at the community level, liable to systematic bias and lacks contextual analyses which recognize that perception from past events can influence responses to future events (Macchi et al. 2014). Perceived climate change and variability analysis is fundamental for gaining a better understanding of the impact of climate change, supplements scientific climate analysis and helps improve the coarse spatial resolution of climate forecasts by using simple practical approximations. However, it has its own limitations in that farmers are unable to remember long-time climate events, inconsistency in use of terms and failure to differentiate climate and weather patterns, has no standard and might be unreliable and focused on agronomic drought which is based on the livelihood impacts the climate has on individual farmers. For instance, even during a year of above average annual rainfall according to observed data, farmers may consider it as a year of agronomic drought ("bara gogiinsa") if their crop failed for some other reasons. This failure might be due to anthropogenic drought that has been brought about by management or policy or institutional failures that reduce or increase the farming system's sensitivity to drought.

7 Conclusion

It is apparent that climate change is a reality at the global and national levels. This study proved that climate change and variability is also a reality at the very local level. The mean maximum and minimum temperature from Arsi Negele Station (7.35°N, 38.66°E, 1913 m a.s.l.) for the period 1983–2014 showed an increasing trend with annual increase of 0.047 and 0.028 °C, respectively. The total rainfall has declined by 10.16 mm per annum. The increase in temperature and decline in total annual rainfall from observed data and reconstructed grid cells had been validated with farmers' perception. Perception of long-term changes in climate by farmers is a fundamental pre-indicator in undertaking adaptation process to calm down the impacts of climate change and variability on farmers' livelihoods and plays an important role in shaping their behaviour. The current situation of climate in the study area, as compared to 30 years ago, the total rainfall perceived to be decreased, the rainfall distribution perceived to be erratic, rainfall onset perceived to be delayed and ended earlier, average temperature perceived to be increased, frost and hail-storm frequencies perceived to be decreased, drought frequency perceived to be increased and flood frequency perceived to be more or less balanced between increase and decrease. Perceptions of risk and cognitive process of primary decision makers are important for the adoption of different adaptation decisions such as adjusting cropping calendar with the shift in the rainfall. This study has come up with that communities have several indigenous knowledges of predicting climatic situations which play important roles in climate risk management. Therefore, it calls for any planned interventions by stakeholders to incorporate with the perception and indigenous knowledge of people to come up with concrete solution for climate change and variability impacts on human livelihoods.

Acknowledgements The authors would like to express their sincere gratitude to the Ethiopian Environment and Forest Research Institute and Wondo Genet College of Forestry and Natural Resources for providing the financial support to Zenebe Mekonnen to carry-out this research as part of his PhD studies. Thanks to the interviewees, the enumerators and Arsi Negele District staffs of the Bureau of Agriculture, especially Aman Gemechu, in contributing their share for the fruitfulness of this research.

Author's contribution ZM contributed to research design, data collection and analysis, manuscript development and review; HK participated in research design and manuscript review; TW contributed to research design and manuscript review; ZA participated in research design and manuscript review.

Compliance with ethical standards

Conflict of interest The authors declare that there is no conflict of interest.

References

- Abid, M., Scheffran, J., Schneider, U. A., & Ashfaq, M. (2014). Farmers' perceptions of and adaptation strategies to climate change and their determinants; the case of Punjab province, Pakistan. *Earth Systems Dynamics*, 5, 1359–1406.
- Adem, A., & Bewket, W. (2011). *A climate change country assessment report for Ethiopia*. Addis Ababa: Epsilon International R&D.
- Amdu, B., Ayehu, A., & Deressa, A. (2013). Farmers' perception and adaptive capacity to climate change and variability in the upper catchment of Blue Nile, Ethiopia. African Technology Policy Studies Network, ATPS Working Paper No. 77. Nairobi, Kenya.
- Amekudzi, L. K., Yamba, E. I., Preko, K., Asare, E. O., Aryee, J., Baidu, M., et al. (2015). Variabilities in rainfall onset, cessation and length of rainy season for the various agro-ecological zones of Ghana. *Climate*, 3, 416–434.

- Asante, F. A., & Amuakwa-Mensah, F. (2015). Climate change and variability in Ghana: Stocktaking. *Climate*, 3, 78–99.
- Bartlett, J. E., Kotrlík, J. W., & Higgins, C. C. (2001). Organizational research: Determining appropriate sample size in survey research. *Information Technology, Learning, and Performance Journal*, 19(1), 43–50.
- Berman, R. J., Quinn, C. H., & Paavola, J. (2014). Identifying drivers of household coping strategies to multiple climatic hazards in Western Uganda: Implications for adapting to future climate change. *Climate and Development*. doi:10.1080/17565529.2014.902355.
- Bernard, H. R. (2006). *Research methods in anthropology: Qualitative and quantitative approaches* (4th ed., p. 803). New York: AltaMira Press.
- Burnham M (2014). The Human Dimensions of Climate Change: Smallholder Perception and Adaptation in the Loess Plateau Region of China. All Graduate Theses and Dissertations. Paper 3331.
- CSA (Central Statistical Agency). (2005). Addis Ababa, Ethiopia.
- Dazé, A., Ambrose, K., & Ehrhart, C. (2009). *Climate vulnerability and capacity analysis handbook*. Atlanta, GA: CARE International.
- Dinku, T., Block, P., Sharoff, J., Hailemariam, K., Osgood, D., del Corral, J., et al. (2014). Bridging critical gaps in climate services and applications in Africa. *Earth Perspectives*, 1(15), 1–13.
- Dinku, T., Hailemariam, K., Maidment, R., Ross, Tarnavsky E., & Connor, S. (2013). Combined use of satellite estimates and rain gauge observations to generate high-quality historical rainfall time series over Ethiopia. *International Journal of Climatology*. doi:10.1002/joc.3855.
- Egeru, A. (2012). Role of Indigenous knowledge in climate change adaptation: A case study of the Teso Sub-Region, Eastern Uganda. *Indian Journal of Traditional Knowledge*, 11(2), 217–224.
- ENC. (2007). Addis Ababa, Ethiopia.
- ENC (Ethiopian National Census). (1994). Addis Ababa, Ethiopia.
- FAO. (2014). Agriculture, forestry and other land use emissions by sources and removals by sinks: 1990–2011 Analysis. Working Paper Series ESS/14- 02. Rome, Italy.
- Ford, J. D., Cameron, L., Rubis, J., Maillet, M., Nakashima, D., Willox, A. C., et al. (2016). Including indigenous knowledge and experience in IPCC assessment reports. *Nature Climate Change*, 6, 349–353.
- Gandure, S., Walker, S., & Botha, J. J. (2013). Farmers' perceptions of adaptation to climate change and water stress in a South African rural community. *Environmental Development*, 5, 39–53.
- Grothmann, T., & Patt, A. (2005). Adaptive capacity and human cognition: the process of individual adaptation to climate change. *Global Environmental Change*, 15, 199–213.
- Gyampoh, B. A., Amisah, S., Idinoba, M., & Nkem, J. (2009). Using traditional knowledge to cope with climate change in rural Ghana. In *Unasylva* 231/232, Vol. 60, pp. 70–74.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., et al. (2012). Carbon emissions from land use and land-cover change. *Biogeosciences*, 9, 5125–5142.
- Huda, M. N., Hossain, M. Z., Ashik-E-Elahi, S., & Mahub, F. (2016). Socio-demographic and economic correlates of climate change coping and adaptation strategies: A study on the farmer communities in Barisal district, Bangladesh. *American Journal of Climate Change*, 5, 167–177.
- IFAD. (2016). The Traditional Knowledge Advantage: Indigenous peoples' knowledge in climate change adaptation and mitigation strategies. International Fund for Agricultural Development (IFAD), Rome, Italy.
- IPCC. (2007). In M. L. Parry, O. F. Canziani, J. P. Palutikof, P. J. van der Linden, & C. E. Hanson (Eds.), *Climate change 2007: Impacts, adaptation and vulnerability. contribution of working group II to the fourth assessment report of the intergovernmental panel on climate change* (p. 976). Cambridge: Cambridge University Press.
- IPCC. (2010a). *Review of the IPCC processes and procedures*, report by the InterAcademy Council (IPCC-32/7), 32nd Session, Bussan, Seoul, 11–14 October 2010. IPCC.
- IPCC. (2013). In T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley (Eds.), *Climate change 2013. The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change* (p. 1535). Cambridge: Cambridge University Press.
- IPCC. (2014). In R. K. Pachauri & L. A. Meyer (Eds.), *Climate change 2014: Synthesis report. Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change* (p. 151). Geneva: IPCC.
- Israel, G. D. (1992). *Determining sample size*. Agricultural Education and Communication Department, University of Florida, IFAS Extension, PEOD6.

- Okonya, J. S., Syndikus, K., & Kroschel, J. (2013). Farmers' perception of and coping strategies to climate change: Evidence from six agro-ecological zones of Uganda. *Journal of Agricultural Science*, 5(8), 252–263.
- Kumar, V. (2014). Role of Indigenous knowledge in climate change adaptation strategies: A study with special reference to North-Western India. *Geography & Natural Disasters*, 5, 131. doi:10.4172/2167-0587.1000131.
- Macchi, M., Gurung, A. M., & Hoermann, B. (2014). Community perceptions and responses to climate variability and change in the Himalayas. *Climate and Development*. doi:10.1080/17565529.2014.966046.
- Magni, G. (2016). Indigenous knowledge and implications for the sustainable development agenda. *Paper commissioned for the Global Education Monitoring Report 2016, Education for people and planet: Creating sustainable futures for all*.
- McLean, K. G., Ramos, C. A., & Rubis, J. (Eds). (2011). Indigenous peoples marginalized populations and climate change: Vulnerability, adaptation and traditional knowledge. *Proceedings of the expert workshop on Indigenous peoples marginalized populations and climate change: Vulnerability, adaptation and traditional knowledge*. Mexico City, Mexico.
- McPhaden, M. J. (2015). Playing hide and seek with El Niño. *Nature Climate Change*, 5, 791–795.
- McSweeney, C., New, M., & Lizcano, G. (2008). UNDP climate change country profiles—Ethiopia. <http://country-profiles.geog.ox.ac.uk>.
- Mekonnen, A., & Hailemariam, K. (2000). *Baseline and climate change scenarios for Ethiopia*. Addis Ababa: NMSA.
- Mertz, O., Mbow, C., Reenberg, A., & Diouf, A. (2009). Farmers' perceptions of climate change and agricultural adaptation strategies in rural Sahel. *Environmental Management*, 43, 804–816.
- Moyo, M., Mvumi, B. M., Kunzekweguta, M., Mazvimavi, K., Craufurd, P., & Dorward, P. (2012). Farmer perceptions on climate change and variability in semi-arid Zimbabwe in relation to climatology evidence. *African Crop Science Journal*, 20(2), 317–335.
- Muluneh, A., Bewket, W., Keesstra, S., & Stroosnijder, L. (2016). Searching for evidence of changes in extreme rainfall indices in the Central Rift Valley of Ethiopia. *Theoretical and Applied Climatology*. doi:10.1007/s00704-016-1739-4.
- NAC (National Academy of Science). (2010). *Adapting to the impacts of climate change* (p. 292). Washington, D.C.: The National Academies Press.
- Nakashima, D. J., McLean, K. G., Thulstrup, H. D., Ramos, C. A., & Rubis, J. T. (2012). *Weathering uncertainty: Traditional knowledge for climate change assessment and adaptation* (p. 120). Paris: UNESCO.
- Ndambiri, H. K., Ritho, C. N., & Mbogoh, S. G. (2014). An evaluation of farmers' perceptions of and adaptation to the effects of climate change in Kenya. *International Journal of Food and Agricultural Economics*, 1(1), 75–96.
- NMA (National Meteorological Agency). (2006). National Adaptation Programme of Action of Ethiopia (NAPA), Addis Ababa.
- NMSA (National Meteorological Service Agency). (1987). Addis Ababa, Ethiopia.
- Nyanga, P. H., Johnsen, F. H., Aune, J. B., & Kalinda, T. H. (2011). Smallholder farmers' perceptions of climate change and conservation agriculture: Evidence from Zambia. *Journal of Sustainable Development*, 4(4), 73–85.
- OFDA/CRED (Office of Foreign Disaster Assistance/Center for Research on the Epidemiology of Disasters). (2009). *Ethiopia country profile, natural disasters*. EM-DAT: The OFDA/CRED International Disaster Database, www.emdat.be—Université catholique de Louvain—Brussels—Belgium.
- Osbah, H., Peter Dorward, P., Stern, R., & Cooper, S. (2011). Supporting agricultural innovation in Uganda to respond to climate risk: Linking climate change and variability with farmer perceptions. *Agricultural Innovation and Climate Risk*, 47(2), 293–316.
- Ramos, C. A., & McLean, K. G. (2012). Climate change mitigation with local communities and indigenous peoples: Practices, lessons learned and prospects. *Proceedings of the international expert workshop climate change Mitigation with Local Communities and Indigenous Peoples held from 26 to 28 March 2012 in Cairns, Austria*. United Nations University Traditional Knowledge Initiative, Darwin, Australia.
- Regassa, S., Givey, C., & Castillo, G. E. (2010). *The rain doesn't come on time any more: poverty, vulnerability, and climate variability in Ethiopia*. Addis Ababa: Oxfam International Research Report.
- Simelton, E., Quinn, C. H., AntwiAgyei, P., Batisani, N., Dougill, A. J., Dyer, J., et al. (2011). *African farmers' perceptions of erratic rainfall*. Centre for Climate Change Economics and Policy, Working Paper No. 73. London, UK.
- Smith, H. A., & Sharp, K. (2012). Indigenous climate knowledges. *Climate Change*, 3, 467–476.

- Stocker, B. D., & Joos, F. (2015). Quantifying differences in land use emission estimates implied by definition discrepancies. *Earth System Dynamics*, 6, 731–744.
- The Anchorage Declaration. (2009). *Indigenous peoples' global summit on climate change, Anchorage Alaska, April 24th 2009*.
- Tierney, J. E., Ummenhofer, C. C., & deMenocal, P. B. (2015). Past and future rainfall in the Horn of Africa. *Climatology*. doi:[10.1126/sciadv.1500682](https://doi.org/10.1126/sciadv.1500682).
- Tilahun, U., & Bedemo, A. (2014). Farmers' perception and adaptation to climate change: Heckman's two stage sample selection model. *Ethiopian Journal of Environmental Studies & Management*, 7, 832–839.
- UNFCCC. (2007). *Climate change: Impacts, vulnerabilities and adaptation in developing countries*. Bonn: Climate Change Secretariat.
- UNFCCC. (2010). *Report of the conference of the parties on its 16th session (FCCC/CP/2010/7/Add.1)*, Cancun, Mexico, 29 November–10 December 2010. Bonn, Germany, UNFCCC.
- Varadan, R. J., & Kumar, P. (2014). Indigenous knowledge about climate change: Validating the perceptions of dryland farmers in Tamil Nadu. *Indian Journal of Traditional Knowledge*, 13(2), 390–397.
- World Bank. (2005). *Ethiopia: Risk and vulnerability assessment*. Report No. 26275-ET. Human Development Group III, Africa Region.
- Yegbemey, R. N., Kabir, H., Awoye, O. H. R., Yabi, J. A., & Paraíso, A. A. (2014). Managing the agricultural calendar as coping mechanism to climate variability: A case study of maize farming in northern Benin, West Africa. *Climate Risk Management*. doi:[10.1016/j.crm.2014.04.001](https://doi.org/10.1016/j.crm.2014.04.001).